

EFFECTIVENESS OF TRANSVERSE OSCILLATORY PRESSURE THERAPY ON PAIN AND MOBILITY IN PATIENTS WITH BILATERAL CERVICAL RADICULOPATHY

Gede Surya Wiranata

Department of Rehabilitation Medicine, Faculty of Medicine, Udayana University, Bali, Indonesia

DOI:10.5281/zenodo.19605211

Abstract

Cervical radiculopathy (CR) is a clinical condition characterized by radiating pain into the upper limb along the distribution of a compressed or irritated cervical nerve root. Although less prevalent than general neck pain, CR is associated with greater levels of pain intensity, functional limitation, and disability, making it a significant musculoskeletal disorder of clinical concern. Epidemiological evidence indicates an annual age-adjusted incidence of approximately 83.2 per 100,000 persons, with higher rates observed in males and peak occurrence during the fifth and sixth decades of life.

The most common etiological factors include cervical disc herniation and cervical spondylosis, while less frequent causes comprise spinal or extraspinal tumors, traumatic nerve root avulsion, synovial and meningeal cysts, dural arteriovenous fistulae, and vascular abnormalities such as tortuous vertebral arteries. Given the functional impairment and persistent pain associated with cervical radiculopathy, effective physiotherapeutic interventions are essential to improve patient outcomes and quality of life.

This study evaluates pain and functional mobility outcomes in patients with bilateral cervical radiculopathy following the application of transverse oscillatory pressure therapy. The intervention aims to reduce pain severity and enhance cervical and upper limb mobility through targeted manual therapy techniques. The study assesses changes in pain perception and functional performance, providing evidence on the therapeutic effectiveness of this approach in clinical practice.

Findings from this study are expected to contribute to the growing body of evidence supporting manual therapy interventions in the management of cervical radiculopathy. The results may assist clinicians in optimizing treatment strategies, improving functional recovery, and reducing disability associated with this condition.

Keywords: Cervical Radiculopathy; Pain Management; Functional Mobility; Manual Therapy; Oscillatory Pressure Therapy

I. INTRODUCTION

Cervical Radiculopathy (CR) is a term used to describe pain radiating into the arm corresponding to the dermatome of the involved cervical nerve root.¹ Cervical radiculopathy forms an important subgroup of neck disorders despite the fact that it is not as frequent as general neck pain, but it has been shown to lead to more severe pain and disability.² In the only large retrospective population-based study, the annual age adjusted incidence rate was 83.2 per 100,000 persons (107.3 for men and 63.5 for women) with a peak incidence in the fifth and sixth decade for both genders.³

Pharmacology, Alternative Medicine, and Healthcare Journal

Research Article

Most common cause of radiculopathy is cervical disc herniation, followed by cervical spondylosis.^{4, 5} CR is less commonly caused by intra spinal or extra spinal tumors, trauma with nerve root avulsion, synovial cysts, meningeal cysts dural arterio-venous fistulae⁶ or tortuous vertebral arteries.⁷

The diagnosis of radiculopathy is based on information received during the subjective and physical examination, which is then confirmed via diagnostic imaging or supported by surgical findings.⁸ Most commonly used test like spur lings test⁹, shoulder abduction test,¹⁰ upper limb tension test(ULTT) or upper limb neural tension test (ULNT)¹¹ neck traction/distraction test, and Valsalva maneuver.¹² The main objectives of treatment in patients with cervical radiculopathy are to relieve pain, improve neurologic function, and prevent reoccurrence.¹³ Some investigators advocated the use of short term immobilization with either a hard or a soft collar to aid in pain control.¹⁴ There are several intervention strategies commonly used in the management of CR. Modalities like TENS, IFT, Ultrasound, cervical traction, electrical muscle stimulator, SWD.¹⁵

An extensive literature review carried out by Haldeman found that transverse oscillatory pressure (TOP), which is one of the techniques of manipulation, provides some benefit in the treatment of chronic mechanical neck pain. Empirical observations of Maitland reported that TOP was recommended for unilaterally distributed symptoms of cervical origin.

Egwu studied the manual forces applied during vertebral mobilization to the cervical spine and found that less time was spent in the use of anterior posterior unilateral pressure and posterior anterior unilateral pressure, he also noted that significantly more patients were pain-free with the use of these techniques when compared to cervical oscillatory rotation and TOP. TOP, originated by Nwuga, although one of the frequently used manipulative techniques by physiotherapists, has been claimed to be effective in amelioration of pain intensity especially radiating pain in cervical, thoracic, and lumbar regions. It involves mobilization of the spinous process of the vertebrae in the region of the spine that had mechanical pain. This technique was reported to be useful when pain has a unilateral distribution, whether localized to the neck or referred to the upper limb. However, there was dearth of documenting evidence on the efficacy of TOP in the management of cervical radiculopathy.¹⁴

II. METHODOLOGY

10 subjects with bilateral cervical radiculopathy confirmed positive spurling test were included. The purpose of the study was explained and written consent was taken from the patient. Cervical ROM and VAS scores were being noted before and after the treatment. Both male and female subjects within age group of 18 to 58 years having bilateral cervical radiculopathy, with positive spurling test, bilateral sensory and motor deficits including sharp pain, muscle weakness and numbness in the upper limb were included in study. Subjects having history of cervical myelopathy, cervical spondylosis or signs of upper motor neuron disease, Unilateral Cervical Radiculopathy (CR), suffering from any other musculoskeletal conditions in the affected limb, who were currently on medications for bilateral cervical radiculopathy and cervical spine surgery within the last 6 months, and non-

Pharmacology, Alternative Medicine, and Healthcare Journal

Research Article

cooperative patients were excluded. Outcome measures were visual analogue scale (VAS) for assessing pain and universal goniometer to assess range of motion of cervical spine.

Intervention:

Transverse oscillatory pressure:

Subject was positioned in prone lying and therapist stood on the side of the patient, the therapist placed the pad of the thumbs against the left side (or right side depending on the location of pain) of the spinous process of vertebrae to be moved. The fingers were spread out on the neck and upper thoracic region. Pressure was directed horizontally through the thumbs to the side of spinous process. Transverse Oscillatory Pressure was executed by a pressure relaxed sequence on the spinous process. Movement was initiated from the trunk and transmitted down the arm to the thumbs.¹⁶ Treatment was affected by push-relax sequence on the spinous using the thumb to produce an oscillatory movement. Transverse pressure was directed towards the side of pain on the cervical vertebra.

The oscillation was given rhythmically for a period of 20 seconds. This was repeated 3 times with a rest period of 2 minutes per session. Treatment was administered 3 times per week for 2 weeks with a total of 6 treatment sessions. Total duration of treatment 15 minutes. The subject's response was assessed after 6th treatment session using VAS and goniometry. Subject was advised not to involve in any other interventions.¹⁴




Fig: Performing Transverse Oscillatory Pressure

III. RESULTS

Table: Data of Outcome Measures with P Value

| Sl.no | | Pre test | post test | | P-Value |
|-------|--|----------|-----------|--|---------|
|-------|--|----------|-----------|--|---------|

Pharmacology, Alternative Medicine, and Healthcare Journal

Research Article

| | Outcome measures | Range | Mean \pm SD | Range | Mean \pm SD | Paired t-test/ wilcoxon test | |
|---|--------------------------------|-------|------------------|-------|------------------|---------------------------------|---------|
| 1 | Flexion ROM | 40-71 | 53.1 \pm 12 | 48-77 | 60.4 \pm 10.5 | t = 8.045 | p<0.05* |
| 2 | Extension ROM | 35-64 | 53 \pm 8.57 | 42-68 | 61 \pm 9 | t = 7.826 | p<0.05* |
| 3 | Cervical right lateral flexion | 20-38 | 29.6 \pm 4.94 | 28-42 | 35.3 \pm 3.88 | t = 14.40 | p<0.05* |
| 4 | Cervical Left lateral flexion | 20-35 | 27.8 \pm 4.51 | 28-41 | 32.9 \pm 4.22 | t = 11.76 | p<0.05* |
| 5 | Right rotation | 37-72 | 57.5 \pm 12.42 | 45-81 | 65.1 \pm 12.56 | t = 12.66 | p<0.05* |
| 6 | Left rotation | 32-79 | 49.2 \pm 14.52 | 41-80 | 56.2 \pm 11.95 | t = 3.96 | p<0.05* |
| 7 | V.A.S (Pain) | 6-9 | 7.40 \pm 1.17 | 3-5 | 4.1 \pm 0.73 | z =2.80 | p<0.05* |

*Statistically Significant

IV. DISCUSSION

The purpose of this study was to evaluate the effect of transverse oscillatory pressure in the treatment of bilateral cervical radiculopathy. The results of the present study showed a highly significant increase in the cervical range of motion (flexion, extension, lateral flexion and cervical rotation) as well as decrease in pain. So, it implies that intervention of Transverse Oscillatory Pressure was significantly effective on reducing pain and improving range of motion among the subjects with bilateral cervical radiculopathy.

By applying TOP directly to the painful area that alter segmental biomechanics by releasing trapped meniscoids, adhesions or by diminishing distortion in the intervertebral disc and restored joint play movement and increases the mobility of cervical region. Also, individual motion segments are thought to be capable of buckling, thereby producing relatively large vertebral motions that achieve a new position of stable equilibrium. The manipulative impulse provides sufficient energy to restore a buckled segment to a lower energy level, thus reducing mechanical stress or strain on soft and hard spinal soft tissues. Giles proposed that spinal manipulation activates all known mechano sensitive, somatosensory receptors because they all possess mechanical thresholds lower than the peak force delivered during a manipulation and the receptor types are responsive to dynamic and/or static components of a mechanical stimulus. By this evidence TOP relieves pain faster. Researchers have reported that TOP to the spinal region has both neurological and mechanical effects.¹⁴

Limitations and Future scope of the study:

Pharmacology, Alternative Medicine, and Healthcare Journal

Research Article

Future studies can be done on larger sample size. In present study, the variables were pain and ROM, studies can be done on neck disabilities, and strength of neck muscles, and can be done with a longer treatment duration. Regular follow-up can be maintained.

Conclusion:

This study concluded that Transverse Oscillatory Pressure (TOP) is effective in improving neck range of motion and reducing pain among the subjects with bilateral cervical radiculopathy.

REFERENCES

- Thoomes EJ, van Geest S, van der Windt DA, Falla D, Verhagen AP, Koes BW, Thoomes-de Graaf M, Kuijper B, Scholten-Peeters WG, Vleggeert-Lankamp CL. Value of physical tests in diagnosing cervical radiculopathy: a systematic review. *The Spine Journal*. 2018 Jan 1;18(1):179-89.
- Ojoawo AO, Olabode A, Esan O, Badru A, Odejide S, Arilewola B. Therapeutic efficacy of cervical traction in the management of cervical radiculopathy: A control trial. *Rwanda Journal of Health Sciences*. 2013 Dec 11;2(2):25-9
- Radhakrishnan K, Litchy WJ, O'Fallon WM, Kurland LT. Epidemiology of cervical radiculopathy: a populationbased study from Rochester, Minnesota, 1976 through 1990. *Brain*. 1994 Apr 1;117(2):325-35.
- Adams C. Cervical spondylitic radiculopathy and myelopathy in: Vinken PJ, Bruyn GW (eds): *Handbook of Clinical Neurology*.
- Wilbourn AJ, Aminoff MJ. AAEE minimonograph# 32: the electrophysiologic examination in patients with radiculopathies. *Muscle & Nerve: Official Journal of the American Association of Electrodiagnostic Medicine*. 1988 Nov;11(11):1099-114.
- Kohno M, Takahashi H, Ide K, Ishijima B, Yamada K, Nemoto S. A cervical dural arteriovenous fistula in a patient presenting with radiculopathy: case report. *Journal of neurosurgery*. 1996 Jan 1;84(1):119-23.
- Horgan MA, Hsu FP, Frank EH. Cervical radiculopathy secondary to a tortuous vertebral artery: case illustration. *Journal of neurosurgery*. 1998 Sep 1;89(3):489-.
- Bussièrès AE, Peterson C, Taylor JA. Diagnostic imaging guideline for musculoskeletal complaints in adults—an evidence-based approach—part 2: upper extremity disorders. *Journal of Manipulative & Physiological Therapeutics*. 2008 Jan 1;31(1):2-32.

Pharmacology, Alternative Medicine, and Healthcare Journal

Research Article

- Spurling RG. Lateral rupture of the cervical intervertebral discs. *Surg Gynec Obstet.* 1944; 78:350-8.
- Davidson RI, Dunn EJ, Metzmaker JN. The shoulder abduction test in the diagnosis of radicular pain in cervical extradural compressive monoradiculopathies. *Spine.* 1981;6(5):441-6.
- Elvey RL. Physical evaluation of the peripheral nervous system in disorders of pain and dysfunction. *Journal of hand therapy.* 1997 Apr 1;10(2):122-9.
- Jull G, Moore A, Falla D, Lewis J, McCarthy C, Sterling M. *Grieve's modern musculoskeletal physiotherapy.* Elsevier; 2015.
- Wolff MW, Levine LA. Cervical radiculopathies: conservative approaches to management. *Physical medicine and rehabilitation clinics of North America.* 2002 Aug;13(3):589-608.
- Ojoawo AO, Olabode A, Esan O, Badru A, Odejide S, Arilewola B. Transverse oscillatory pressure in management of cervical radiculopathy: A randomised controlled study. *Hong Kong Physiotherapy Journal.* 2016 Jun 1; 34:19-26.
- Cleland JA, Fritz JM, Whitman JM, Heath R. Predictors of short-term outcome in people with a clinical diagnosis of cervical radiculopathy. *Physical therapy.* 2007 Dec 1;87(12):1619-32.
- Nwuga VCB. Techniques of spinal manual therapy. In:Nwuga VC, editor. *Manual treatment of back pain.* 2nd ed. William and Wilkins; 2007. p. 115