

QUANTITATIVE ANALYSIS OF DRUG DISTRIBUTION USING COMPARTMENTAL MODELS

Dr. Fatima Hauwa Bello

Department of Mathematics and Statistics, Federal University Otuoke, Bayelsa State, Nigeria

DOI:10.5281/zenodo.19663638

Abstract

Alcohol addiction poses serious health and social challenges, making the study of ethanol metabolism in the human body an important area of research. Most existing studies on ethanol kinetics focus on single-dose experiments designed to determine the rate of alcohol elimination or to assess impairment, particularly in relation to highway safety. However, real-life drinking patterns often involve repeated alcohol consumption, which introduces additional complexity in modeling ethanol absorption, distribution, and elimination.

Repeated dosing scenarios require more sophisticated analytical approaches, and as a result, many studies have avoided such investigations. Mathematical modeling, particularly compartmental models, provides an effective framework for analyzing these complex dynamics. Compartmental models describe the movement of substances between different physiological compartments and help capture the processes involved in ethanol metabolism.

This study applies a compartmental mathematical approach to analyze the kinetics of ethanol metabolism in the human body. The model represents the body as interconnected compartments through which ethanol is absorbed, distributed, and eliminated over time. The approach provides insight into the behavior of alcohol concentration under different intake conditions and highlights the usefulness of mathematical models in understanding alcohol metabolism.

Keywords: Alcohol metabolism, Compartmental modeling, Ethanol kinetics, Mathematical modeling, Blood alcohol concentration

INTRODUCTION

Alcohol addiction is a phenomenon that has attracted the attention of several researchers and academics from a wide range of professions due to the devastating effects it has on many facets of human existence, Tireito (2022). Ethanol metabolism has primarily been based on single-dose experimental investigations to map the time course of ethanol elimination or from the perspective of highway safety according to Saha et al., (2015). When alcohol is used in repeated doses, further problems occur, Rota et al. (2014).

The model becomes more sophisticated and requires more analysis when several ingestions are simulated. As a result, both experimentalists and modelers have simply avoided multiple-dose investigations in favour of single-dose research Rota *et al.* (2014). Mathematical models, more specifically compartment models that examine the fluxes between the compartments, may handle such research rather readily, Plawewcki et al., (2008). The majority of the compartment models that currently exist for alcohol removal are deterministic. Although such models offer a wealth of knowledge regarding the pharmacokinetics of ethanol, their effectiveness is hindered by the need to determine the model parameters, such as the anatomical structure of the assumed compartments, metabolic rates, as well as transient and steady state solutions. Stochastic models, which offer relevant information with little

complexity, are a front-runner for modelling such phenomena due to the variability in the instants of alcohol consumption, absorption, and elimination times, which are random in nature. Based on the above mentioned challenges, there are other researchers who have venture into pharmacokinetics modelling. Shirley (2007) employed Markov or hidden Markov models in one study to analyze respondents' drinking patterns. Instead of being steady across time, these models are better suited to depict systems that undergo abrupt shifts. Wang et al. (2002) multivariate temporal models for alcohol intake were recommended in another investigation. Guang (1998) provided a stochastic model using queuing theory in another study to understand the harmful effects of ethanol on the body.

For the ethanol elimination process, a stochastic model was put up by Ghadirinejad *et al.* (2016). Their model was notable for accounting for numerous doses, zero and first order elimination, and other factors. Their findings were able to take into consideration changes in the quantity and timing of alcohol consumption. According to Plawecki *et al.* (2008), human-based compartment models that result in differential equation systems are typical mathematical models for blood alcohol content. Each variable in these models represents the amount of ethanol present in a particular organ or system, such as the vascular or digestive systems. A differential equation can be used to characterize the rate at which ethanol is absorbed, transported to another compartment, or metabolized. Models range in complexity from simple single-compartment models, Umulis *et al.* (2005) to more complex multi-compartment models (Ramchandani *et al.*, 1999; Pieters *et al.*, 1990) with three compartments or more. Bunonyo and Amadi (2023) employed system of mathematical model to analyze the concentration of alcohol in the GI tract and bloodstream. Their models were solved analytically and some of the useful entry parameters were obtained. According to their findings, alcohol content in a compartment rises as the pace of alcohol movement in that compartment increases. Smith et al. (1993) looked at the rate at which ethanol left the circulation in healthy males and females as well as in alcohol abusers. Following the oral administration of ethanol (0.8 g/kg body weight) in the form of whiskey, blood alcohol levels were estimated over the course of three hours. In Meem (2022) employed a mathematical model for alcohol issues, there are four demographic classes represented by his model: potential drinkers, occasional drinkers, heavy drinkers, and recovered and abstainers. Sensitivity analysis of the basic reproduction number shows that focusing solely on alcoholics is less effective in the long run at preventing the spread of alcoholism than encouraging and supporting potential drinkers to refrain from adopting drinking habits and occasional drinkers to stop drinking. A mathematical model of alcohol misuse created by Sharma and Samanta (2015) has four compartments that correspond to the four demographic classes, moderate, and infrequent drinkers, heavy drinkers, drinkers under treatment, and temporarily recovered class. To comprehend the dispersion of medication administration in the human body via oral and intravenous routes, Khanday *et al.* (2017) created mathematical models, using Fick's principle and the law of mass action, three models were created based on the diffusion process. To find the answer to the ordinary differential equations relating to the rate of change of concentration in various compartments, including blood and tissue medium, the

Laplace transform and eigenvalues methods were applied. According to the findings, the drug concentration gradually rises in the other compartments while falling in the first.

Rockerbie and Rockerbie (1995) developed a mathematical model using an iterative algorithm based on the Widmark formula and used it to simulate a continuous spectrum of the expected blood alcohol concentration from start of alcohol dosage to time of specific event in numerical and graphic form. The 2-compartment model employed by Norberg et al. (2000) consists of the peripheral compartment, which houses the remaining body fluids in other tissues, and the central compartment, which in this case contains the blood plasma and the tissues that are in quick equilibrium with it (liver and kidney). When the alcohol concentration dropped to levels about 0.10 g/l and lower, they demonstrated that the rate of alcohol elimination transitioned to a linear phase on their semi-logarithmic.

Using a mathematical model, Hongzhi (2023) investigated how alcohol diffuses throughout the body. Under the fundamental presumption that the body's process for transferring alcohol is separated into four stages (alcohol-stomach-body-fluids-body), a mathematical model is developed, and numerical simulation is used to verify the diffusion process when consuming in various ways. Pieters *et al.* (1990) used a three-compartment model to describe the pharmacokinetics of alcohol following oral delivery.

Moore (2022) forecast blood ethanol concentration levels over time for a variety of drinking and eating scenarios by included the process of food digestion in the model. Four categories of drinkers were defined by Baker *et al.* (2014): very heavy drinkers, heavy drinkers, binge drinkers, and light drinkers. The average daily intake of very heavy drinkers (VHD) is more than 3 g/kg, and more than 10% of their daily intakes are higher than 4 g/kg. Heavy drinkers (HD) continue to consume more than 3 g/kg per day on more than 20% of their drinking days. However, this study is an extension of Bunonyo *et al.* (2023), whose investigation is limited to the drug concentration in the body and bloodstream compartments without consideration of how the elimination and metabolic rate affect the gastrointestinal and urinary tracts. In this research we shall adopt the Laplace method in solving the models we would be formulating.

MATERIALS AND METHODS

This study investigates the administration of drugs and the concentration of drug from one compartment to another, as seen in Figure 1. The diffusion of drugs from one compartment to another is with specific rate of diffusion because of the complexity of different human body system. The compartment diagram helps us with the formulation of the first order ordinary differential equations that mimics the diffusion and concentration of drugs from the gastrointestinal tract, the body, the bloodstream and the urinary tracts respectively.

Drug in Bloodstream




Figure1 Diagram showing drug diffusion from one compartment to another with different rate

Assumptions

Before we can go ahead and formulate the system of mathematical models, let's consider the following assumptions:

- The concentration of drugs in the body compartment can be affected by the increasing rate of inter-compartmental distribution.
- The concentration of drugs in the bloodstream compartment is affected by the intercompartmental distribution rates.
- The elimination rate affects the drug concentration in the body and bloodstream compartments.
- The first-order rate constant affects the drug concentration in the body and bloodstream compartments.
- The metabolic rate constant affects the drug concentration in the body and bloodstream compartments.
- The volume and dosage of the drug contribute to its concentration in the body and bloodstream compartments.

Definition of Parameters c_1 : Concentration of drug in the human body compartment c_2 : Concentration of drug in bloodstream compartment

c_u : Concentration of drug in urinary tract c_{GI} : Concentration of drug in gastrointestinal tract k_{12}, k_{21} : Inter-compartmental distribution rate constant k_e : Excretion rate constant k_m : Metabolic rate constant k_a : First order rate constant

V : Volume of drug in the compartments

D : Drug dosage in the compartments

Research Article

Model Formulation

The formation of the system of mathematical models of the problem under investigation follows the above compartment diagram in Figure 1, Bunonyo *et al.* (2023) and Bunonyo and Amadi (2023), where the system of models is formulated as:

$$\frac{dc_1}{dt} = kac_{GI} - kac_1 - k_{21}c_2 - k_{12}c_1 - kec_1 - kmc_1 - kec_u \tag{1}$$

$$\frac{dc_2}{dt} = k_{12}c_1 - k_{21}c_2 \tag{2}$$

$$\frac{dc_u}{dt} = kec_1 - kec_u \tag{3}$$

$$\frac{dc_{GI}}{dt} = ka c_{GI} \tag{4}$$

Equations (1)-(4) are subject to certain initial conditions, they are: $c_1(0) = 0, c_2(0) = 0,$

$$c_u(0) = 0, c_{GI}(0) = 0$$

$$D = \begin{pmatrix} -kac_1 - k_{21}c_2 - k_{12}c_1 - kec_1 - kmc_1 - kec_u & 0 & 0 & 0 \\ k_{12}c_1 & -k_{21}c_2 & 0 & 0 \\ kec_1 & -kec_u & 0 & 0 \\ ka c_{GI} & 0 & 0 & 0 \end{pmatrix} \tag{5}$$

Methods of Solution

Considering the concentration of the administered drug via the GI on other compartments, we have to solve equation (4) first, that is:

$$\frac{dc_{GI}}{dt} = ka c_{GI}$$

Recalling that the Laplace method can be stated as:

$$L\{c_{GI}(t)\} = c_{GI}(s) = \frac{ka}{s} c_{GI}(t) \tag{7}$$

Applying equation (7) on equation (6), we have:

$$L\{dc_{GI}\} = ka L\{c_{GI}\} \tag{8}$$

$$s c_{GI} - c_{GI}(0) = ka c_{GI} \tag{9}$$

Simplifying equation (9) with the initial condition in equation (5), we have:

$$D c_{GI} = \frac{ka}{s} c_{GI} \tag{10}$$

Research Article

Simplifying equation (16), we have the following:

$$D c_{GI}(s) + V(s)k_a = \dots \quad (11)$$

Taking the inverse of the Laplace transform in equation (11), we have:

$$c_{GI}(t) = L^{-1}\{c_{GI}(s)\} \quad (12)$$

Applying equation (12) in solving equation (11), we have: $c(t) = L^{-1}\{c(s)\}$

□

$$\frac{dc_{GI}}{dt} = k c \quad (6)$$

$$s^2 k_{21}$$

Solving the model representing the drug concentration in the body, we apply the Laplace method on equation (1), which is:

$$s c_1 + k_{21} c_1 - k_{12} c_2 = k_a c_1 - k_e c_1 - k_m c_1 - k_e c_u \quad (26)$$

Substituting equations (17), (26) and (30) into equation (33), we have:

$$k_a L c_{GI} - k_a L c_1 - k_{21} L c_2 - k_{12} L c_1 - k_e L c_1 - k_m L c_1 - k_e L c_u = 0 \quad (24)$$

Simplifying equation (31), we have:

$$k_a L c_{GI} - k_a L c_1 - k_{21} L c_2 - k_{12} L c_1 - k_e L c_1 - k_m L c_1 - k_e L c_u = 0 \quad (27)$$

Simplifying equation (27), we have:

$$k_a L c_{GI} - k_a L c_1 - k_{21} L c_2 - k_{12} L c_1 - k_e L c_1 - k_m L c_1 - k_e L c_u = 0 \quad (25)$$

$$k_a L c_{GI} - k_a L c_1 - k_{21} L c_2 - k_{12} L c_1 - k_e L c_1 - k_m L c_1 - k_e L c_u = 0 \quad (28)$$

where $k_{12} = k_a + k_e + k_m$

Simplifying equation (28), we have:

$$k_a L c_{GI} - k_a L c_1 - k_{21} L c_2 - k_{12} L c_1 - k_e L c_1 - k_m L c_1 - k_e L c_u = 0 \quad (29)$$

Simplifying equation (29), we have:

$$k_a L c_{GI} - k_a L c_1 - k_{21} L c_2 - k_{12} L c_1 - k_e L c_1 - k_m L c_1 - k_e L c_u = 0 \quad (30)$$

Simplifying equation (30), we have:

$$k_a L c_{GI} - k_a L c_1 - k_{21} L c_2 - k_{12} L c_1 - k_e L c_1 - k_m L c_1 - k_e L c_u = 0 \quad (31)$$

Research Article

$$L^1 = c_1 s^3 + c_2 s^2 + c_3 s + c_4 \quad (36)$$

where $k_1, k_2, k_3, k_4, k_5, k_6, k_7, k_8, k_9, k_{10}, k_{11}, k_{12}, k_{13}, k_{14}, k_{15}, k_{16}, k_{17}, k_{18}, k_{19}, k_{20}, k_{21}, k_{22}, k_{23}, k_{24}, k_{25}, k_{26}, k_{27}, k_{28}, k_{29}, k_{30}, k_{31}, k_{32}, k_{33}, k_{34}, k_{35}, k_{36}, k_{37}, k_{38}, k_{39}, k_{40}, k_{41}, k_{42}, k_{43}, k_{44}, k_{45}, k_{46}, k_{47}, k_{48}, k_{49}, k_{50}, k_{51}, k_{52}, k_{53}, k_{54}, k_{55}, k_{56}, k_{57}, k_{58}, k_{59}, k_{60}, k_{61}, k_{62}, k_{63}, k_{64}, k_{65}, k_{66}, k_{67}, k_{68}, k_{69}, k_{70}, k_{71}, k_{72}, k_{73}, k_{74}, k_{75}, k_{76}, k_{77}, k_{78}, k_{79}, k_{80}, k_{81}, k_{82}, k_{83}, k_{84}, k_{85}, k_{86}, k_{87}, k_{88}, k_{89}, k_{90}, k_{91}, k_{92}, k_{93}, k_{94}, k_{95}, k_{96}, k_{97}, k_{98}, k_{99}, k_{100}$

$$L^2 = A s^3 + B s^2 + C s + D \quad (37)$$

If we express $s^3 = k_1 s^2 + k_2 s + k_3$, then equation (36) becomes:

$$L^1 = c_1 (k_1 s^2 + k_2 s + k_3) + c_2 s^2 + c_3 s + c_4 \quad (38)$$

Simplifying equation (38), we have:

$$s^2 (k_1 c_1 + c_2) + s (k_2 c_1 + c_3) + (k_3 c_1 + c_4) = A s^3 + B s^2 + C s + D \quad (39)$$

Substituting equation (39) into equation (538), we have:

$$L^1 = c_1 (k_1 s^2 + k_2 s + k_3) + c_2 s^2 + c_3 s + c_4 = A s^3 + B s^2 + C s + D \quad (40)$$

Simplifying equation (39), we have:

$$s^2 (k_1 c_1 + c_2) + s (k_2 c_1 + c_3) + (k_3 c_1 + c_4) = A s^3 + B s^2 + C s + D \quad (41)$$

Simplifying equation (41), we have: $A s^3 + B s^2 + C s + D = s^2 (k_1 c_1 + c_2) + s (k_2 c_1 + c_3) + (k_3 c_1 + c_4)$

$$s^2 (k_1 c_1 + c_2) + s (k_2 c_1 + c_3) + (k_3 c_1 + c_4) = A s^3 + B s^2 + C s + D \quad (42)$$

Upon further simplification of equation (42), we have:

$$2 k_1 s^2 + k_2 s + k_3 = A s^3 + B s^2 + C s + D \quad (43)$$

From equation (43), we have:

Research Article

$$A \square B \square C \square D \square 0 \tag{44}$$

$$A \square r_3 \square r_2 \square r_1 \square B \square r_3 \square k_a \square r_2 \square C \square r_1 \square k_a \square r_3 \square D \square r_1 \square k_a \square r_2 \square 1 \tag{45}$$

$$A \square r r_2 \square r_3 \square r r_1 \square r_3 \square r r_1 \square 2 \square B \square r k_3 \square a \square r r_3 \square 2 \square r k_2 \square a \square C \square r k_1 \square a \square r r_1 \square 3 \square r k_3 \square a \square D \square r k_1 \square a \square r r_1 \square 2 \square r k_2 \square a \square k \tag{46}$$

$$A \square r r_1 \square 2 \square 3 \square B \square r r k_2 \square 3 \square a \square C \square r r k_3 \square 1 \square a \square D \square r r k_2 \square 1 \square a \square k_{21} \square k_e \tag{47}$$

Equations (44)-(47) can be expressed as:

$$\begin{aligned} & \square \quad 1 \quad 1 \quad 1 \quad 1 \quad \square \square A \square \square 0 \square \\ & \square \square \square \square r_2 r_3 \square r r_1 \square 3 \square r r_1 \square 2 \square \square k_a \square r_2 \square r_3 \square \quad \square k_a \quad 3 \square \quad \square k_a \square r_1 \square r_2 \square \quad \square \square B \square \square \square 1 \square \square \\ & \square r_1 \square r \quad \square \square \square \\ & \square \quad \square \square \square \square \quad \square \\ & \square \quad \square \square \square \square \quad \square \\ & \square \square r_2 r_3 \square r r_1 \square 3 \square r r_1 \square 2 \square \quad \square r_3 r_2 \square r_3 k_a \square r_2 k_a \square \square r r_1 \square 3 \square r k_1 \square a \square r_3 k_a \square \square r_2 k_a \square r k_1 \square a \square r r_1 \square 2 \square \square \square C \square \square \\ & k \square \square \\ & \square \quad \square \square \square \square \quad \square \\ & \square \square \square \square r r_1 \square 2 r_3 \quad r_2 r_3 \quad \square \square \square \square D \square \square \square \square \square \square \square k_{21} \square k_e \square \square \square k_a \quad r_3 r k_1 \square a r_2 r k_1 \square a \square \square \end{aligned} \tag{48}$$

Further simplification of equation (48), we have:

$$\begin{aligned} & \square 1 \quad 1 \quad 1 \quad 1 \quad \square \square A \square \square 0 \square \\ & \square r \square r \square 1 \quad r \square 1 \quad r \square 1 \quad r \square 1 \quad \square \square \square \square B \square \square \square \square 1 \quad \square \square \\ & \square \quad 1 \\ & \square \quad \square \square \square \square \quad \square \\ & \square \quad \square \square \square \square \quad \square \tag{49} \\ & \square r \square 2 \square r \square 2 \quad r \square 2 \quad r \square 2 \quad \square \square C \square \square k \square \square \\ & \square \quad \square \square \square \square \quad \square \end{aligned}$$

$$\square \square \square \square r r r_1 \square 2 \square 3 \square r r k_2 \square 3 \square a \square r r k_3 \square 1 \square a \square r r k_2 \square 1 \square a \square \square \square \square \square \square D \square \square \square \square \square \square \square k_{21} \square k_e \square \square \square \text{ where } r_{\square 1} \square \square \square r_2 r_3 \square r r_1 \square 3 \square r r_1 \square 2 \square, r_{\square 1} \square \square \square k_a \square r_2 \square r_3 \square, r_{\square 1} \square \square \square k_a \square r_1 \square r_3 \square, r_{\square 1} \square \square \square k_a \square r_1 \square r_2 \square, r_{\square 2} \square \square \square r r_2 \square 3 \square r r_1 \square 3 \square r r_1 \square 2 \square, r_{\square 2} \square \square \square r r_3 \square 2 \square r k_3 \square a \square r k_2 \square a \square, r_{\square 2} \square \square \square r r_1 \square 3 \square r k_1 \square a \square r k_3 \square a \square, r_{\square 2} \square \square \square r k_2 \square a \square r k_1 \square a \square r r_1 \square 2 \square$$

Recall that $\square s \square k \square s \square k \square s \square k \square \square \square \square s \square r_1 \square \square s \square r_2 \square \square s \square r_3 \square \tag{50}$

Solving equation (50), where r_1, r_2 and r_3 denotes the roots of the equation (50). To obtain the roots, we adopt Wolfram Mathematica, which are:

Research Article

$$r_1 = \frac{k_\beta}{3} + \frac{2^{1/3}(k_\beta^2 - 3k_\gamma)}{3 \left(-2k_\beta^3 + 9k_\beta k_\gamma - 27k_\phi + \sqrt{-4(k_\beta^2 - 3k_\gamma)^3 + (2k_\beta^3 - 9k_\beta k_\gamma + 27k_\phi)^2} \right)^{1/3}} + \frac{\left(-2k_\beta^3 + 9k_\beta k_\gamma - 27k_\phi + \sqrt{-4(k_\beta^2 - 3k_\gamma)^3 + (2k_\beta^3 - 9k_\beta k_\gamma + 27k_\phi)^2} \right)^{1/3}}{3 \times 2^{1/3}} \tag{51}$$

$$r_2 = -\frac{k_\beta}{3} + \frac{(1 + \sqrt[3]{3})(-k_\beta^2 + 3k_\gamma)}{3 \times 2^{2/3} \left(-2k_\beta^3 + 9k_\beta k_\gamma - 27k_\phi + \sqrt{-4(k_\beta^2 - 3k_\gamma)^3 + (2k_\beta^3 - 9k_\beta k_\gamma + 27k_\phi)^2} \right)^{1/3}} + \frac{(-1 + \sqrt[3]{3}) \left(-2k_\beta^3 + 9k_\beta k_\gamma - 27k_\phi + \sqrt{-4(k_\beta^2 - 3k_\gamma)^3 + (2k_\beta^3 - 9k_\beta k_\gamma + 27k_\phi)^2} \right)^{1/3}}{6 \times 2^{1/3}} \tag{52}$$

$$r_3 = s + \frac{k_\beta}{3} + \frac{(1 + \sqrt[3]{3}) \left(-2k_\beta^3 + 9k_\beta k_\gamma - 27k_\phi + \sqrt{-4(k_\beta^2 - 3k_\gamma)^3 + (2k_\beta^3 - 9k_\beta k_\gamma + 27k_\phi)^2} \right)^{1/3}}{6 \times 2^{1/3}} + \frac{(1 - \sqrt[3]{3})(-k_\beta^2 + 3k_\gamma)}{3 \times 2^{2/3} \left(-2k_\beta^3 + 9k_\beta k_\gamma - 27k_\phi + \sqrt{-4(k_\beta^2 - 3k_\gamma)^3 + (2k_\beta^3 - 9k_\beta k_\gamma + 27k_\phi)^2} \right)^{1/3}} \tag{53}$$

where A, B, C and D are constants and can be obtained by solving equation (49)

Simplifying equation (40), we obtained:

$$c1 \square t \square \square L \square 1 \square c1 \square s \square \square \square \square 0AL \square 1 \square \square \square \square 1 \square \square \square \square 0BL \square 1 \square \square \square 1 \square \square \square \square 0CL \square 1 \square \square \square 1 \square \square \square \square 0DL \square 1 \square \square \square \square \square \square (54)$$

$$\square \square \square s \square ka \square \square \square \square \square s \square r_1 \square \square \square s \square r_2 \square \square \square s \square r_3 \square$$

Finally, upon adopting the basic Laplace technique, equation can be simplified as:

$$c1 \square t \square \square L \square 1 \square c1 \square s \square \square \square \square 0Ae \square kat \square \square 0Bert1 \square \square 0Cer2t \square \square 0Der3t \tag{55}$$

RESULTS

The numerical simulation of the analytical solutions of the system of mathematical models formulated in our previous sections was done using Wolfram Mathematica, version 12, and the graphical results are presented as follows:




Figure 2 Effect of elimination rate on drug concentration in the body and the urinary tract compartments

Research Article




Figure 3 Effect of metabolic rate on drug concentration in the body compartment




Figure 4 Effect of first order rate constant on drug concentration in the gastrointestinal tract compartment




Figure 5 Effect of first order rate constant on drug concentration in the body and bloodstream compartments




Figure 6 Effect of the inter-compartment distribution rate on drug concentration in the body and bloodstream compartments




Figure 7 Effect of the inter-compartment distribution rate on drug concentration in the body and bloodstream compartments

DISCUSSION

In Figure 2, we noticed that the drug concentration in the body increases for the elimination rate at 0.5 unit furthermore, the drug concentration in the body keeps decreases for an increase in the elimination rate from 0.5 to 1.5 units. The result also showed that the drug concentration decreases for time for as fast as the rate become. The rate of metabolism effect on drug concentration in the body was also investigated and found out that for the metabolic rate at 0.3 unit, the drug concentration in the body decrease after attaining different maximum height as seen in Figure 3.

Figure 4 depicts the effect of the first rate constant on drug concentration in gastrointestinal tract compartment. It can be seen that the rate constant of 0.7 per hour, the drug concentration in the gastrointestinal tract decreases, for an increase in rate constant, furtherance of the investigation, we can see that the gastrointestinal drug concentration continuously decreased to zero stability level at the different first order rate constant. Figure 5 illustrates the influence of the first rate constant on drug concentration in the body and in the bloodstream compartments for different rate constants. It can be seen that the rate constant of 0.7 unit per hour, the drug concentration in the body and in the bloodstream compartment increases, however, the investigation revealed that the drug concentration in both compartments increased to a maximum before decreasing to steady state over time.

Research Article

Figures 6 and 7 illustrate the influence of the inter-compartmental rates of the drug concentration in the bloodstream and body compartments. The results indicated that as the inter-compartmental rate k_{12} term increases the drug concentration levels in the body compartment decrease, however, it has been observed that the increase k_{12} term the drug concentration in the bloodstream compartment increases.

CONCLUSION

In conclusion, we have been able to formulate a system of mathematical models to mimic the diffusion of drug and the concentration of drug from one compartment to another where the first compartment denotes the gastrointestinal, the second the body, the fourth is the bloodstream, and finally, the urinary tract compartments respectively. In a similar vein, we developed computational codes using Wolfram Mathematica to simulate the effect of the various rates on the drug concentration in the body and bloodstream compartments as part of the research objectives, and based on research investigation, we conclude as follows:

- The drug concentration in the body keeps decreases for an increase in the elimination rate from 0.5 to 1.5 units. This showed that for one to quickly elimination the drug concentration in the body system, we must accelerate the elimination rate.
- The rate of metabolism affects the drug concentration in the body with a little increase in metabolic rate.
- To decrease the drug concentration in the gastrointestinal tract, there should be an increase in rate constant.
- In order to reduce the drug concentration in the body and the bloodstream compartments, one has to increase the rate constant.
- The inter-compartmental rates increase influenced the drug concentration in the bloodstream and body compartments.

REFERENCE

- Tireito, F., Apima, S., & Muchika, C. L. (2022). Mathematical Modelling of Alcoholism Incorporating Media Awareness as an Intervention Strategy.
- Saha, D., Alluri, P., & Gan, A. (2015). Prioritizing Highway Safety Manual's crash prediction variables using boosted regression trees.
- Rota, M., Porta, L., Pelucchi, C., Negri, E., Bagnardi, V., Bellocco, R., ... & La Vecchia, C. (2014). Alcohol drinking and multiple myeloma risk—a systematic review and meta-analysis of the dose–risk relationship.
- Plawecki, M. H., Han, J. J., Doerschuk, P. C., Ramchandani, V. A., & O'Connor, S. J. (2008). Physiologically based pharmacokinetic (PBPK) models for ethanol.
- Shirley, K. E. (2007). *Hidden Markov models for alcoholism treatment trial data*. University of Pennsylvania

Research Article

- Wang, S. J., Winchell, C. J., McCormick, C. G., Nevius, S. E., & O'Neill, R. T. (2002). Short of complete abstinence: an analysis exploration of multiple drinking episodes in alcoholism treatment trials. *Alcoholism*:
- Guang, W. (1998). Application of queueing theory with Monte Carlo simulation to the study of the intake and adverse effects of ethanol.
- Ghadirinejad, M., Atasoylu, E., Izbirak, G., & Matina, G. S. (2016). A stochastic model for the ethanol pharmacokinetics.
- Umulis, D. M., Gürmen, N. M., Singh, P., & Fogler, H. S. (2005). A physiologically based model for ethanol and acetaldehyde metabolism in human beings. *Alcohol*, 35(1), 3-12.
- Ramchandani, V. A., Bolane, J., Li, T. K., & O'Connor, S. (1999). A physiologically-based pharmacokinetic (PBPK) model for alcohol facilitates rapid BrAC clamping.
- Pieters, J. E., Wedel, M., & Schaafsma, G. (1990). Parameter estimation in a three-compartment model for blood alcohol curves.
- Bunonyo, K. W., & Amadi, U. C. (2023). Alcohol Concentration Modelling in the GI Tract and the Bloodstream.
- Smith, G. D., Shaw, L. J., Maini, P. K., Ward, R. J., Peters, T. J., & Murray, J. D. (1993). Mathematical modelling of ethanol metabolism in normal subjects and chronic alcohol misusers.
- Meem, I. J., Hossain, R., & Samad, S. A. (2022). A mathematical model of alcoholism in bangladesh.
- Sharma, S., & Samanta, G. P. (2015). Analysis of a drinking epidemic model.
- Khanday, M. A., Rafiq, A., & Nazir, K. (2017). Mathematical models for drug diffusion through the compartments of blood and tissue medium.
- Rockerbie, D. W., & Rockerbie, R. A. (1995). Computer simulation analysis of blood alcohol.
- Norberg, Å., Gabrielsson, J., Jones, A., & Hahn, R. G. (2000). Within-and between-subject variations in pharmacokinetic parameters of ethanol by analysis of breath, venous blood and urine.
- Hongzhi, Z. (2023). Mathematical model and numerical simulation on the diffusion of alcohol in the human body.

Research Article

- Moore, S., Radunskaya, A., Zollinger, E., Grant, K. A., Gonzales, S., Walter, N. A., & Baker, E. J. (2022). Pairing food and drink: A physiological model of blood ethanol levels for a variety of drinking behaviors.
- Baker, E. J., Farro, J., Gonzales, S., Helms, C., & Grant, K. A. (2014). Chronic alcohol self-administration in monkeys shows long-term quantity/frequency categorical stability.
- Bunonyo, K. W., Ebiwareme, L., & Okrinya, A. B. (2023). Mathematical Modeling of Alcohol Drug Concentration in the Bloodstream.
- Bunonyo, K. W., Ebiwareme, L., & Awomi, P. Z. (2023). Temperature effect on drug diffusion in the stomach and bloodstream compartments.